Statistical Theory and Modeling (ST2601) Lecture 10 - Logistic, Poisson regression and Beyond

Mattias Villani

Department of Statistics Stockholm University













Linear Gaussian regression as a probability model

Logistic regression

Poisson regression

Generalized linear models and beyond

Linear Gaussian Regression

The usual formulation for the *i*th observation

$$Y_i = \beta_0 + \beta_1 x_{1i} + \beta_2 x_{2i} + \ldots + \beta_p x_{pi} + \varepsilon_i, \quad \varepsilon_i \stackrel{\text{iid}}{\sim} N(0, \sigma_{\varepsilon}^2)$$

The usual formulation in vector form

$$Y_i = \mathbf{x}_i^\top \boldsymbol{\beta} + \varepsilon_i, \quad \varepsilon_i \stackrel{\text{iid}}{\sim} N(0, \sigma_{\varepsilon}^2)$$

Equivalent formulation

$$\begin{aligned} \mathbf{Y}_i \mid \mathbf{x}_i \overset{\text{ind}}{\sim} \mathbf{N}(\mu_i, \sigma_{\varepsilon}^2) \\ \mu_i = \mathbf{x}_i^{\top} \boldsymbol{\beta} \end{aligned}$$

Regression is a model for a conditional distribution $f(y|\mathbf{x})$.

The *i*th observation has its own mean μ_i given by the regression line.

Regression models a conditional distribution Y|x



Logistic regression for binary response variable

Assume now that the response Y_i is binary (0 or 1). Without covariates: model distribution as Bernoulli $Y_i \stackrel{\text{iid}}{\sim} \text{Bernoulli}(\mu)$

With covariates: model **conditional distribution** as Bernoulli $Y_i \mid \mathbf{x}_i \stackrel{\text{ind}}{\sim} \text{Bernoulli}(\mu_i)$

Modeling the conditional mean as $\mu_i = \mathbf{x}_i^\top \boldsymbol{\beta}$ is no good.

Use logistic function $f(z) = \frac{1}{1+e^{-z}}$ to ensure that $0 \le \mu_i \le 1$

$$\mu_i = \Pr(Y_i = 1 \mid \boldsymbol{x}_i) = \frac{1}{1 + e^{-\boldsymbol{x}_i^\top \boldsymbol{\beta}}}$$

Logistic regression

$$Y_i \mid \boldsymbol{x}_i \stackrel{\text{ind}}{\sim} \text{Bernoulli}(\mu_i)$$
$$\mu_i = \frac{1}{1 + e^{-\boldsymbol{x}_i^\top \boldsymbol{\beta}}}$$

Maximum likelihood for Bernoulli data

■ Model: Y₁,..., Y_n ~ Bernoulli(p) [Note µ = 𝔼(Y_i) = p]
 ■ Bernoulli probability function

$$p(y) = \begin{cases} 1 - p & \text{if } y = 0\\ p & \text{if } y = 1 \end{cases}$$

or

$$\boldsymbol{p}(\boldsymbol{y}) = \boldsymbol{p}^{\boldsymbol{y}}(1-\boldsymbol{p})^{1-\boldsymbol{y}}$$

Likelihood

$$\prod_{i=1}^{n} p(y_i|p) = \prod_{i=1}^{n} p^{y_i} (1-p)^{1-y_i} = p^{\sum_{i=1}^{n} y_i} (1-p)^{n-\sum_{i=1}^{n} y_i}$$
$$= p^s (1-p)^f$$

■ $s = \sum_{i=1}^{n} y_i$ is the number of successes ■ f = n - s is the number of failures.

Maximum likelihood for Bernoulli data

Likelihood

$$L(p) = p^{s}(1-p)^{f}$$

Log-likelihood

$$\ell(p) = s \log p + f \log(1-p)$$

First derivative (recall: $f(x) = \log(x)$ then f'(x) = 1/x)

$$\ell'(p) = \frac{s}{p} - \frac{f}{1-p}$$

Maximum likelihood estimate \hat{p} is the *p* that solves

$$\ell'(p) = \frac{s}{p} - \frac{f}{1-p} = 0$$

which has solution

$$\hat{p} = \frac{s}{n}$$

Maximum likelihood for Bernoulli data - widget



Maximum likelihood for Logistic regression 🕯

Logistic regression

$$Y_i \mid \boldsymbol{x}_i \stackrel{\text{ind}}{\sim} \text{Bernoulli}(\mu_i)$$
$$\mu_i = \frac{1}{1 + e^{-\boldsymbol{x}_i^\top \boldsymbol{\beta}}}$$

- **Data**: responses \mathbf{y} ($\mathbf{n} \times 1$) and covariates $\mathbf{X}(\mathbf{n} \times \mathbf{p})$.
- Likelihood function (covariates assumed fixed, non-random

$$\begin{split} L(\boldsymbol{\beta}) &= \prod_{i=1}^{n} p(y_i | \boldsymbol{x}_i) \\ &= \prod_{i=1}^{n} \mu_i^{y_i} (1 - \mu_i)^{1 - y_i} \\ &= \prod_{i=1}^{n} \left(\frac{1}{1 + e^{-\boldsymbol{x}_i^\top \boldsymbol{\beta}}} \right)^{y_i} \left(\frac{e^{-\boldsymbol{x}_i^\top \boldsymbol{\beta}}}{1 + e^{-\boldsymbol{x}_i^\top \boldsymbol{\beta}}} \right)^{1 - y_i} \end{split}$$

Numerical maximization with optim.

Poisson regression for count data

Assume now that the response Y_i is a count (0, 1, 2, ...).

Without covariates: **distribution** is Poisson

 $Y_i \stackrel{\text{iid}}{\sim} \operatorname{Poisson}(\mu)$

With covariates: conditional distribution is Poisson

$$Y_i \mid \mathbf{x}_i \stackrel{\text{ind}}{\sim} \operatorname{Poisson}(\mu_i)$$

Modeling the conditional mean as $\mu_i = \mathbf{x}_i^\top \boldsymbol{\beta}$ is no good.

Use exponential function to ensure that $\mu_i > 0$

$$\mu_i = e^{\mathbf{x}_i^\top \boldsymbol{\beta}}$$

Poisson regression

$$Y_i \mid \boldsymbol{x}_i \stackrel{\text{ind}}{\sim} \text{Poisson}(\mu_i)$$
$$\mu_i = \mathbb{E}(Y_i \mid \boldsymbol{x}_i) = e^{\boldsymbol{x}_i^\top \boldsymbol{\beta}}$$

ML for Poisson regression - widget



Generalized linear models (GLM)

Continuous positive data. Gamma regression

$$Y_i \mid \boldsymbol{x}_i \stackrel{\text{ind}}{\sim} \text{Gamma}(\alpha, \boldsymbol{e}^{\boldsymbol{x}_i^\top \boldsymbol{\beta}})$$
$$\mu_i = \mathbb{E}(Y_i \mid \boldsymbol{x}_i) = \alpha \boldsymbol{e}^{\boldsymbol{x}_i^\top \boldsymbol{\beta}}$$

- Data as proportions: **Beta regression**.
- **Truncated data**: truncated normal regression: widget
- ... and so on ...
- Generalized Linear Models.
 - Maximum likelihood by numerical maximization.
- Sampling distribution from Observed information

$$\hat{\boldsymbol{eta}} \stackrel{\mathrm{approx}}{\sim} \mathsf{N}\left(\boldsymbol{eta}, \mathcal{J}_{\mathsf{n}}^{-1}(\hat{\boldsymbol{eta}})
ight)$$
 for large n

GLMs are linear models

Logistic regression

$$Y_i \mid \mathbf{x}_i \stackrel{\text{ind}}{\sim} \text{Bernoulli}(\mu_i)$$
$$\mu_i = \frac{1}{1 + e^{-\mathbf{x}_i^\top \boldsymbol{\beta}}}$$

Decision boundary: $Pr(y_i = 1 | \mathbf{x}_i) = Pr(y_i = 0 | \mathbf{x}_i)$

$$\mu_{i} = 1 - \mu_{i} \iff \frac{1}{1 + e^{-\mathbf{x}_{i}^{\top}\boldsymbol{\beta}}} = \frac{e^{-\mathbf{x}_{i}^{\top}\boldsymbol{\beta}}}{1 + e^{-\mathbf{x}_{i}^{\top}\boldsymbol{\beta}}}$$
$$1 = e^{-\mathbf{x}_{i}^{\top}\boldsymbol{\beta}} \iff 0 = \mathbf{x}_{i}^{\top}\boldsymbol{\beta}$$

(take log on both sides, and recall $\log 1 = 0$ and $\log e^a = a$).

Decision boundaries are linear in the features, x.
 Linear GLMs are:

 highly interpretable.

- robust to overfitting.
- restrictive.

Logistic regression - linear decision boundaries





5-nearest neighbour





Non-linear regression

Example: Poisson non-linear regression

$$Y_i \mid \mathbf{x}_i \stackrel{\text{ind}}{\sim} \operatorname{Poisson}(\mu_i)$$

 $\mu_i = e^{f(\mathbf{x}_i)}$

where f(x_i) is some (non-linear) function of the covariates.Examples:

► Linear:
$$f(x) = \beta_0 + \beta_1 x$$

► Polynomial: $f(x) = \beta_0 + \beta_1 x + \beta_2 x^2 + \beta_3 x^3 + \dots$

Other non-linear models:

Splines

- Regression trees
- Neural networks
- Gaussian processes